

Living and Dying on the Streets: Palliative Care for Structurally Vulnerable Populations

Dr. Trevor Morey, MD, CCFP(PC), Palliative Care Physician

Leeann Trevors, MSW, Patient Navigator



Land Acknowledgment




Source: CSC Data Warehouse. (June 2018).

- •Indigenous peoples now account for 27% of the total federal inmate population. 37.6% of the federal women inmate population is Indigenous.
- •Between 2007 and 2016, the Indigenous population has increased by 39%, compared to 5% for the overall prison population.
- Despite faster entry into correctional programs and higher program completion rates, Indigenous offenders are still being released later and revoked more often than their counterparts.














- 1. Review the impact of homelessness on health outcomes.
- 2. Review promising interventions to address the supportive and palliative care needs people experiencing homelessness.
- 3. Inspire equity-based directions to improve care for structurally vulnerable populations.







Social determinants of health

Digging at the roots, not just low hanging fruit:

The reproduction of the social determinants of health when the structural determinants' are left untouched

~Dr Nanky Rai





Examples of structural vulnerabilities

- Homeless & vulnerably housed
- Homelessness
- Poverty
- Social isolation
- Substance use
- Trauma
- Poverty


- Social isolation & trauma
- Racialized communities
- Substance use disorders
- Mental illness
- New Canadians, immigrants, refugees
- Non-status populations

Homelessness is a continuum



Living at risk of
homelessness due to lack
of financial security or
other factors (intimate
partner violence,
separation, divorce)

Living in accommodations without security of tenure (couch-surfing, rooming houses)



Quantifying Canada's homeless

- 35 000 people nightly
- 150 000 sheltered annually
- 150 000-300 000 homeless people per year




Did you know?



person in a shelter











The health of people who are homeless

- 75% with one or more chronic disease
 - HCV: 28x
 - Heart Disease: 5x
 - Cancer: 4x

- Acute care:
 - ED: 8x
 - Hospital admission: 4x
- The elderly



The health of people who are homeless

- Highest all-cause mortality rate in Canada
 - Life expectancies:

34 - 47 years old

- Mortality rates: 2.3x 4x
- Location at FOL



50%



Best Practices: Palliative Care

The NEW ENGLAND IOURNAL of MEDICINE

SOUNDING BOARD

Early Specialty Palliative Care — Translating Data in Oncology into Practice

Ravi B. Parikh, A.B., Rebecca A. Kirch, J.D., Thomas J. Smith, M.D., and Jennifer S. Temel, M.D.

Early palliative care for patients with advanced cancer: a cluster-randomised controlled trial



Camilla Zimmermann, Nadia Swami, Monika Krzyzanowska, Breffni Hannon, Natasha Leighl, Amit Oza, Malcolm Moore, Anne Rydall, Gary Rodin, Ian Tannock, Allan Donner, Christopher Lo

Summary

Background Patients with advanced cancer have reduced quality of life, which tends to worsen towards the end of life.

We assessed the effect of early palliative care in patients with advanced cancer on several aspects of quality of life.

Published Online February 19, 2014 http://dx.doi.org/10.1016 S0140-6736/13162416-2

Effects of a Palliative Care Intervention on Clinical Outcomes in Patients With Advanced Cancer

The Project ENABLE II Randomized Controlled Trial

Marie Bakitas, DNSc, APRN	Contex
Kathleen Doyle Lyons, ScD, OTR	tive care
Mark T. Hegel, PhD	Object
Stefan Balan, MD	sympton
rances C. Brokaw, MD, MS	 Design Novemb
anette Seville PhD	- Novemi

Jav G. Hull, PhD

Context There are few randomized controlled trials on the effectiveness of palliative care interventions to improve the care of patients with advanced cancer.

Objective To determine the effect of a nursing-led intervention on quality of life, symptom intensity, mood, and resource use in patients with advanced cancer.

Design, Setting, and Participants Randomized controlled trial conducted from November 2003 through May 2008 of 322 patients with advanced cancer in a rural, National Cancer Institute–designated comprehensive cancer center in New Hampshire and affiliated outreach clinics and a VA medical center in Vermont.

The NEW ENGLAND JOURNAL of MEDICINE

ORIGINAL ARTICLE

Early Palliative Care for Patients with Metastatic Non–Small-Cell Lung Cancer

Jennifer S. Temel, M.D., Joseph A. Greer, Ph.D., Alona Muzikansky, M.A., Emily R. Gallagher, R.N., Sonal Admane, M.B., B.S., M.P.H., Vicki A. Jackson, M.D., M.P.H., Constance M. Dahlin, A.P.N., Craig D. Blinderman, M.D., Juliet Jacobsen, M.D., William F. Pirl, M.D., M.P.H., J. Andrew Billings, M.D., and Thomas J. Lynch, M.D.



Best Practices: Homeless Health

- Outreach
- Intensive case management
- Interdisciplinary
- Care across settings
- Harm reduction
- Integration with housing sector



A new model of care









The PEACH team







About PEACH

- How is the PEACH program structured?
- How do we get referrals?
- Who are our clients?
- What do we do for them (and their communities)?
- What are their trajectories?
- What has PEACH accomplished?



About PEACH

- Reconnection to family or friends
- Prevention of acute hospitalizations
 /ED use
- EOL in preferred place
- Housing status

Table 1: Housing status of PEACH clients at time of referral and time of death

	Time of referral (% of clients)	Time of death (% of clients)		
Shelter	24 (38.1%)	5 (7.9%)		
Affordable Rental	17 (27.0%)	0		
Transitional Housing	12 (19.0%)	8 (12.7%)		
Social Housing	6 (9.5%)	1 (1.6%)		
Sleeping rough	2 (3.2%)	0		
Market Rental	1 (1.6%)	0		
Unknown	1 (1.6%)	6 (9.5%)		
PCU/Hospice	0	28 (44.4%)		
Acute Care Hospital	0	15 (23.8%)		



ICHA | Inner City | Health Associates

59 Adelaide Street East. 2ndfloor, M5C 1K6 Phone: 416-591-4411 Fax: 416-640-2072

EXTERNAL REFERRAL

PALLIATIVE EDUCATION AND CARE FOR THE HOMELESS (PEACH)

The Palliative Education and Care for the Homeless initiative with Inner City Health Associates has been developed to provide palliative supports to individuals with life limiting illnesses in addition to having experienced homelessness or are vulnerably housed.

PEACH Referral Criteria

- o Homeless or vulnerably housed
- Life-threatening illness
- Significant pain and/or symptomatic burden
- Difficulty accessing mainstream palliative care services

PEACH is able to accept a limited number of clients into the program. Please refer to ICHA website for additional Palliative Care supports in the GTA.

We will review the referral and connect with you regarding the referral status.

Thank you



59 Adelaide Street East, 2nd floor, M5C 1K6 Phone: 416-591-4411 Fax: 416-640-2072

EXTERNAL REFERRAL

Palliative Education and Care for the Homeless (PEACH)

	Health Card No.		Version
Referral Source:			
Name	T	itle	Phone Number
Referring Physician	Organization Name		Fax Number
Patient Info:			/ /
First Name	Last Name	Gender	Date of Birth
Phone Number	Address	Address She	
Housing Situation			
Reasons for Referral:			
Reasons for Referral:			
Primary Palliative Diagnosis: Substance Use (type, amount, freque	ency):	Date of Dia	agnosis://
Primary Palliative Diagnosis:	ency):	Date of Dia	agnosis://
Primary Palliative Diagnosis: Substance Use (type, amount, freque Mental Health Diagnosis:	ency):	Date of Dia	ngnosis://
Primary Palliative Diagnosis: Substance Use (type, amount, freque Mental Health Diagnosis: Please note when the referral is ac	ency):	Date of Dia	ngnosis://
Primary Palliative Diagnosis: Substance Use (type, amount, freque Mental Health Diagnosis: List of Wedications List of Medications	ency):	Date of Dia	ngnosis://
Primary Palliative Diagnosis: Substance Use (type, amount, freque Mental Health Diagnosis: Please note when the referral is ac Use of Medications Recent Laboratory results	ency):	Date of Dia	ngnosis://
Primary Palliative Diagnosis: Substance Use (type, amount, freque Mental Health Diagnosis: Please note when the referral is ac iss of Medications Recent Laboratory results Pathology Reports	ency):	Date of Dia	ngnosis://
Primary Palliative Diagnosis: Substance Use (type, amount, freque Mental Health Diagnosis: Please note when the referral is ac Use of Medications Recent Laboratory results	ency):	Date of Dia	ngnosis://
Primary Palliative Diagnosis: Substance Use (type, amount, freque Mental Health Diagnosis: Please note when the referral is ac Use of Medication Recent Laboratory results Pathology Reports Recent Consultation Notes	ency):	Date of Dia	ngnosis://_ will be requested:

Why PEACH works

- Community ← → hospital
- Integration within model of home & community care
- Person-centered (not physician-centered)
- Focused on coordination & health navigation
- Communication without borders
- A community of practice centered on advocacy



On Calgary streets: Dignity, at the end of life



British Columbia

Doctor and nurse go mobile to provide palliative care to Victoria's homeless



Judgment and stigma discourage people from accessing proper care, says local researcher







SDoH & the palliative care GAP


Palliative care patients living in the poorest neighbourhoods (still housed) in Ontario:

- Were least likely to get a home visit from a doctor (29.4% vs 40.2%)
- Were more likely to have unplanned ED visits (65.4% vs 59.8%)
- Were more likely to get admitted to hospital in their last 30 days of life (64.5 % vs 58.9%)

A key determinant of palliative care access



Social Accountability



Micro: The clinical environment; encompasses both the individual family physician—patient relationship and the inter-professional, team-based care setting.



 Meso: The local community; the geographic context in which clinical and academic medical work are situated. Includes education, training, and continuing professional development (CPD).



Macro: The broader realm of policies and their impact on population and public health, where family physicians act as advocates for healthy public policy.





Our Social Accountability

A socially accountable health system is one that dedicates resources to where they are truly needed, up and downstream



Structural vulnerability

"An individual's or a population group's condition of being at risk for negative health outcomes through their interface with socioeconomic, political, and cultural/normative hierarchies. Patients are structurally vulnerable when their location in their society's multiple overlapping and mutually reinforcing power hierarchies (e.g., socioeconomic, racial, cultural) and institutional and policy-level statuses (e.g., immigration status, labor force participation) constrain their ability to access health care and pursue healthy lifestyles."

Death is a social justice issue

People living with life-limiting illness



Affected by deficits in the social determinants



Death is a Social Justice Issue

Advances in Nursing Science
Vol. 39, No. 4, pp. 293–307
Copyright © 2016 Wolters Kluwer Health, Inc. All rights reserved.

Death Is a Social Justice Issue Perspectives on Equity-Informed Palliative Care

Sheryl Reimer-Kirkham, PhD, RN; Kelli Stajduhar, PhD, RN; Bernie Pauly, PhD, RN; Melissa Giesbrecht, PhD; Ashley Mollison, MA; Ryan McNeil, PhD; Bruce Wallace, PhD

All too often, palliative care services are not responsive to the needs of those who are doubly vulnerable, being that they are both in need of palliative care services and experiencing deficits in the social determinants of health that result in complex, intersecting health and social concerns. In this article, we argue for a reorientation of palliative care to explicitly integrate the premises of health equity. We articulate the philosophical, theoretical, and empirical scaffolding required for equity-informed palliative care and draw on a current study to illustrate such an approach to the care of people who experience structural vulnerabilities. **Key words:** discrimination, bealth equity, bomelessness, marginalization, palliative care, poverty, public bealth, social justice, stigma, structural vulnerability


Aging and Dying in Prison

AGING AND DYING IN PRISON

An Investigation into the Experiences of Older Individuals in Federal Custody



Aging and Dying in Prison



Aging and Dying in Prison

Compassionate Release

Two cases were brought forward by CSC staff members during interviews that demonstrate how difficult it is for a terminally ill patient to be allowed to die in the community. In the first case, an elderly patient was dying of cancer in a medium security institution and CSC staff put his case forward to the Parole Board to be granted release to a hospice within the community. The response from the Parole Board was that it wanted to see the offender managed in a minimum security institution before granting him release to the hospice. He was eventually released to the community, but died 2 hours after release.

In the second case, CSC staff supported the transfer of a terminally ill patient (minimum security) to a hospice facility within the community, and brought the case forward to the Parole Board. There was concern that the patient may not have much time left and that they may lose the spot they had secured with the hospice so the institution tried to push the Parole Board to make a decision quickly. The Board responded that should it not be able to make a decision quickly, CSC could use a medical escorted temporary absence (ETA) to accommodate the offender in the community hospice. A medical ETA requires uniformed guards to be present with the offender at all times. Quite appropriately, the hospice was not supportive of these conditions.

Haley House





Report: Too little too late





inews Homeless, vulnerable only find best health care when at death's door, UVic study finds

Homeless, vulnerable only find best health care when at death's door, UVic study finds







Researchers followed 25 marginally housed people for 2 years



NOVEMBER 2018

Thank You!

Dr. Trevor Morey trevor.morey@unityhealth.to

Leeann Trevors trevorsL@smh.ca

Links

Global	News-	PEACH	program
--------	-------	--------------	---------

Part 1:

https://globalnews.ca/news/7478848/coronavirus-toronto-palliative-care-peach-program/

Part 2:

https://globalnews.ca/news/7478961/coronavirus-toronto-peach-program/

Part 3

 $\underline{https://globalnews.ca/video/7489780/good-wishes-project-helps-fulfill-dreams-for-toronto-palliative-care-patients-experiencing-homelessness and the results of the following of the following project of the following pr$

Toronto Star-

Haley House

https://www.youtube.com/watch?v=jl7zMVojUa0

Links

PEACH program

http://www.icha-toronto.ca/programs/peach-palliative-education-and-care-for-the-homeless

University of Victoria Too Little Too Late Report

https://www.uvic.ca/research/groups/peol/assets/docs/too-little-too-late.pdf




Top 10 practices to improve care for structurally vulnerable populations






Derive equity by design









Practice trauma-informed healthcare

Structurally vulnerable populations to consider:

- Homeless & vulnerably housed
- Indigenous communities
- Poverty
- Social isolation & trauma
- Racialized communities

- Substance use disorders
- Mental illness
- New Canadians, immigrants, refugees
- Non-status populations



Operationalize anti-oppressive frameworks



Digging at the roots, not just low hanging fruit:

The reproduction of the social determinants of health when the structural determinants' are left untouched



Decolonize our healthcare systems





Take a human rights approach



Meet people where they are at



Prioritize dignity





Empower the populations we aim to serve



Support each other



Thank you @NightShiftMD and @CBCWhiteCoat for sharing our experience in grieving as a way to prevent burnout. What experiences or rituals can you share that work for you and your teams? #cdnhealth #sdoh #hpm



Palliative care team helps the homeless die 'with dignity,' a healing circle hel... After a homeless woman died of breast cancer, the doctors, nurses and shelter workers who helped to ease her final days gathered to remember her. $\mathscr S$ cbc.ca



Challenge the status quo within our communities

